

SIMPLE WASHING TECHNIQUE FOR REMOVAL OF PESTICIDES AND HEAVY METALS FROM VEGETABLES AND MUSSELS

NURAIN BINTI ABD JALIL

A thesis submitted in fulfilment of the
requirements for the award of the degree of
Master of Science (Chemistry)

Faculty of Science
Universiti Teknologi Malaysia

NOVEMBER 2016

*To my beloved mother and father, En. Abd Jalil Bin Lebai Mat and Pn. Rossnani
Binti Suboh and to my supportive siblings and special appreciation to Assoc. Prof Dr.
Razali Ismail as supervisor.*

ACKNOWLEDGMENT

First of all, thanks to Allah S.W.T for his mercy and guidance in giving I full strength to complete this thesis. Even facing with some difficulties in completing this task, I still managed to complete it. A lot of thanks to my supervisor, Assoc. Prof. Dr. Razali Bin Ismail for guided me in my project over the past years. He has provided me with the knowledge, the tools, and the motivation that I will need to succeed in the future.

Special appreciation to very helpful lab assistants, Madam Norzubaidha Ismail and Ms Siti Nurul Aini Asbullah for giving me chance handling instruments in their lab such as Gas Chromatography with Electron Capture Detector and Inductively Couple Plasma-Optical Electron Spectroscopy. Then, I would like thanks to my parents, for supporting me mentally and physically not just during finishing this tasks but also during my whole studies. Last but not least, grateful acknowledgement to all of my friends who never give up in giving their support to me in all aspects.

Thank you.

ABSTRACT

Washing is a common practice at home and industry to avoid contamination of germs and dangerous diseases and to remove residual contaminants (pesticides and heavy metals) from vegetables. A commercial vegetable washing solution was tested for the removal of nine organophosphate pesticides (dichlorvos, ethoprophos, disulfoton, parathion-methyl, fenclorophos, chlorpyrifos, prothiofos, azinphos-methyl and malathion) by soaking the vegetables in the washing solution. Meanwhile, in the heavy metals removal study, the samples were washed with sodium acetate chelating solution to remove four heavy metals, namely arsenic, lead, cadmium and nickel. Method validation study was carried out for the analysis of pesticides and heavy metal residues in term of linear range, repeatability, recovery, limits of detection and limits of quantification. The optimum washing practices for pesticides removal were achieved with five minute washing time, 15 mL washing solution and at temperature of 27°C with swirling. The optimized conditions for heavy metals removal were 1 hour chelating time at temperature of 29.5°C and sodium acetate concentration of 500 mg/L. Results showed that six of the pesticides residues were within the safe permitted levels except for ethoprophos (1.4 - 0.15 ppm) and disulfoton (0.25 - 0.03 ppm). Meanwhile, the removal effectiveness for pesticides and heavy metals from the real samples were in the range of 16.7% - 97.3% and 5.5% - 100.0%, respectively. Among the heavy metals studied, only nickel (0.16 ppm) in long beans achieved the safe permitted level after washing with sodium acetate solution. The results also suggested that continuous process is a better technique compared to combined or individual process since the continuous process achieved lower residual contaminant levels than the allowed maximum residue limit for certain pesticides and heavy metals.

ABSTRAK

Membasuh adalah amalan biasa di rumah dan industri untuk mengelakkan pencemaran kuman dan penyakit berbahaya dan untuk penyingkiran bahan cemar (sisa racun perosak dan logam berat) dari sayur-sayuran. Sejenis produk larutan basuhan sayur-sayuran komersil telah diuji untuk meningkatkan kadar penyingkiran sembilan racun perosak organofosfat (diklorvos, etoprofos, disulfoton, paration-metil, fenklorfos, klorpirifos, protiofos, azinfos-metill dan malation) dengan merendam sayur-sayuran di dalam larutan basuhan. Sementara itu, dalam kajian penyingkiran logam berat, sampel telah dibasuh dengan larutan pengkelat natrium asetat untuk menyingkirkan empat logam berat iaitu arsenik, plumbum, kadmium dan nikel. Kajian pengesanan kaedah telah dijalankan untuk analisis sisa racun perosak dan logam berat dalam bentuk julat linear, kebolehulangan, perolehan, had pengesanan dan had kuantifikasi. Amalan basuhan yang optimum dicapai bagi penyingkiran sisa racun perosak dengan rendaman selama lima minit, menggunakan 15 mL larutan basuhan dan pada suhu 27°C dengan bantuan pusran. Keadaan optimum bagi penyingkiran logam berat adalah 1 jam bagi masa pengkelatan pada suhu 29.5°C dan kepekatan natrium asetat 500 mg/L. Keputusan menunjukkan bahawa enam jenis sisa racun perosak adalah pada tahap yang dibenarkan kecuali etoprofos (1.4 - 0.15 ppm) dan disulfoton (0.25 - 0.03 ppm). Sementara itu, keberkesanan penyingkiran racun perosak dan logam berat daripada sampel sebenar adalah dalam julat masing-masing di antara 16.7% - 97.3% dan 5.5% - 100.0%. Di antara logam berat yang dikaji, hanya nikel (0.16 ppm) di dalam kacang panjang mencapai tahap selamat yang dibenarkan selepas dibasuh dengan larutan natrium asetat. Keputusan juga mencadangkan bahawa proses berterusan adalah teknik yang lebih baik berbanding proses gabungan atau individu kerana proses berterusan mencapai tahap pencemar baki yang lebih rendah daripada had baki maksimum yang dibenarkan untuk racun perosak dan logam berat yang tertentu.

TABLE OF CONTENTS

CHAPTER	TITLE	PAGE
	DECLARATION	ii
	DEDICATION	iii
	ACKNOWLEDGEMENT	iv
	ABSTRACT	v
	ABSTRAK	vi
	LIST OF TABLES	xii
	LIST OF FIGURES	xv
	LIST OF ABBREVIATIONS	xvii
1	INTRODUCTION	1
	1.1 Background of Study	1
	1.2 Problem of Statement	5
	1.3 Objectives of Study	7
	1.4 Scope of Study	8
	1.5 Significant of Study	9
2	LITERATURE REVIEW	10
	2.1 Worldwide Awareness on Pesticide	10
	2.1.1 Campaign and Organization Involvement	12

2.1.2	Common Pesticides Reported in Vegetable and Mussel	13
2.1.3	Guideline on Maximum Residue Limits	15
2.1.4	Extraction and Determination of Pesticides	16
2.2	Heavy Metals	17
2.2.1	Biological Magnification of Heavy Metal	20
2.2.2	Extraction and Determination of Heavy Metal	21
2.3	Removal Techniques for Pesticide Residue	22
2.4	Washing Technique for Pesticide's Removal	23
2.4.1	Homemade and Commercial Washing Solution	24
2.4.2	Mechanism of Washing Process	25
2.5	Combination of Washing and Chelating Treatment	26
2.5.1	Solutions used in Treatments for Removal of Pesticide Residue	26
2.5.2	Proposed Design for Reduction of Pesticide and Heavy Metal Residues using Continuous Washing Process	28
3	EXPERIMENTAL	29
3.1	Apparatus	29
3.2	Materials	29

3.3	Sampling	30
3.4	Validation Studies	30
3.4.1	Linearity	31
3.4.2	Recovery	32
3.4.3	Repeatability	32
3.4.4	Limit of Detection	34
3.4.5	Limit of Quantification	34
3.5	Screening of Functional Groups in Commercial Washing Solution	35
3.5.1	Sampling and Coating of Vegetables with Pesticides	35
3.5.2	Washing Parameters	35
3.5.3	Extraction of Vegetable Samples	36
3.5.4	Analysis of Pesticides	36
3.6	Chelating Treatment for Heavy Metals	37
3.6.1	Preparation of Chelating Agents	37
3.6.2	Treatment Parameters of Vegetable Samples	38
3.6.3	Digestion of Vegetable Samples	38
3.6.4	Analysis of Heavy Metals	39
3.7	Process of Combination Washing and Chelating Treatment	39
3.7.1	Combine Process	39

	3.7.1.1	Extraction and Analysis of Samples	39
	3.7.2	Continuous Process	40
4		RESULTS AND DISCUSSIONS	42
	4.1	Validation Studies for Pesticides Removal	42
	4.1.1	Analysis of Organophosphorus Pesticides	42
	4.1.2	Calibration Curve (linearity)	43
	4.1.3	Repeatability and Recovery	45
	4.1.4	Limits of Detection (LODs) and Limits of Quantification (LOQs)	48
	4.2	Validation Studies for Heavy Metals Removal	50
	4.2.1	Calibration Curve (linearity)	51
	4.2.2	Repeatability and Recovery	52
	4.2.3	Limits of Detection (LODs) and Limits of Quantification (LOQs)	53
	4.3	Parameters of Washing Treatment	53
	4.3.1	FTIR Spectrum of Commercial Washing Gel	54
	4.3.2	pH of Washing Solution	57
	4.3.3	Physical Changes on Samples	58
	4.3.4	The Normal Practices in Washing Treatment	59

4.3.5	Real Sample Analysis	63
4.4	Evaluation of Heavy Metals Analysis	66
4.4.1	The Optimized Chelating Treatments Conditions	66
4.4.2	Application Chelating Treatment to Vegetable Samples	69
4.5	Application of Washing Technique and Chelating Treatment	72
4.5.1	Combined Process	73
4.5.2	Continuous Process	76
4.6	Mechanism of Washing Solution and Sodium Acetate Chelating Solution	76
5	CONCLUSION AND RECCOMENDATION	80
5.1	Conclusion	80
5.2	Recommendation	82
	REFERENCES	83

LIST OF TABLES

TABLE NO.	TITLE	PAGE
1.1	Review on percentage of residue dissipation	6
2.1	Physical properties of selected pesticides	11
2.2	The percentage of pesticide residues in several samples from year 2000 – 2008	14
2.3	Toxicity of trace heavy metals	19
2.4	Maximum permitted proportion in foods	19
2.5	Percentage reduction of pesticide residues using various washing solutions.	27
4.1	Summary of retention time for investigated analytes	43
4.2	Linearity range, r^2 and equation values of investigated pesticides in hexane:acetone (9:1) solvent	45

4.3	Repeatability and recoveries in fortified vegetables	47
4.4	Comparisons of LODs and LOQs with MRL of the investigated pesticides.	50
4.5	Series of working standards and r^2 of the calibration curves	51
4.6	The results of repeatability and recovery test for mussel and vegetable sample	52
4.7	LODs and LOQs of heavy metals studied.	53
4.8	Peaks on the FTIR spectrum of concentrated washing gel	56
4.9	pH of washing solution	57
4.10	Initial concentration in fortified samples and percentage reduction after washing process	64
4.11	The optimized conditions for removal of heavy metals from vegetable sample	69
4.12	Results of comparison of heavy metal concentrations in some vegetables with WHO/FAO permissible values (WHO, 1999)	70
4.13	Concentration of heavy metals after washed with chelating solution	72

4.14 (a)	The detection and percentage removal of pesticide residues in combined process.	74
4.14 (b)	The detection and percentage removal of heavy metal residues in combined process.	74
4.15 (a)	The detection and percentage removal of pesticide residues in continuous process.	76
4.15 (b)	The detection and percentage removal of heavy metal residues in continuous process.	76

LIST OF FIGURES

FIGURE NO.	TITLE	PAGE
2.1	Molecular structure for studied pesticides	10
3.1	Summary of experimental design for combined process	40
3.2	Summary of experimental design for combined process	41
4.1	Separation peaks (A-J) of 1.00 ppm standard mixture OPP's in hexane:acetone solvent (9:1)	44
4.2	Calibration curves from (a) to (g) were standard from mixture organophosphorus pesticides standard solution	46
4.3	GC-ECD chromatograms obtained for the analysis of broccoli sample (a); for sample before spiking (b); and after spiking with standard mixture OPP's at level of 1.00 ppm	49
4.4	Calibration curves for studied heavy metals (As, Cd, Ni and Pb)	51

4.5	FTIR Spectrum of concentrated washing gel	54
4.6	Difference in appearance washing using tap water (left) and commercial washing solution (right)	58
4.7	Trend of residues reduction with different of washing time	60
4.8	Trend of residues reduction with different in volume of washing solution	61
4.9	Removal rate of residues reduction for different washing temperature.	62
4.10	Removal rate of residues reduction for swirl and unswirl effect	63
4.11	Effect of time in chelating treatment	67
4.12	Effect of concentration in chelating treatment	67
4.13	Effect of temperature in chelating treatment	68
4.14	Percentage removal of heavy metals using chelating agent solution.	71

LIST OF ABBREVIATIONS

ACGIH	-	Association Advancing Occupational and Environmental Health
AR	-	Acid Reduction
As	-	Arsenic
Cd	-	Cadmium
CP	-	Chlorpyrifos
DDE	-	1,1-Dichloro-2,2-bis(p-chlorophenyl) ethylene
DDT	-	Dichlorodiphenyltrichloroethane
DOA		Department of Agriculture
<i>E. coli</i>	-	<i>Escherichia coli</i>
EFLE	-	Easily and Freely Leachable Fraction
EPA	-	Environmental Protection Agency
FDA	-	Food and Drugs Administrative
GC-ECD		Gas Chromatography - Electron Capture Detector
GC/MS	-	Gas Chromatography/ Mass Spectrometer
Hg	-	Mercury
HPLC-DAD	-	High Performance Liquid Chromatography - Diode-Array Detection
HUS	-	Hemolytic Uremic Syndrome

LC-MS/MS	-	Liquid Chromatography – Mass Spectrometer / Mass Spectrometer
LLE	-	Liquid-liquid Extraction
LODs	-	Limit of Detection
LOQs	-	Limit of Quantification
MCLs	-	Maximum Contaminant Levels
MENGO		Malaysian Environmental NGOs
MRLs	-	Maximum Residue Levels
NIOSH	-	National Institute for Occupational Safety and Health
OO	-	Organic Oxidation
OPP	-	Organophosphorus Pesticide
PAN A	-	Pesticide Action Network Asia and the Pacific
Pb	-	Lead
POPs	-	Persistent Organic Pollutants
RR	-	Resistant fraction
SPE	-	Solid Phase Extraction
SPME	-	Solid Phase Micro Extraction
TWA	-	Time-Weighted Average
U. S.	-	United States
USDA	-	United States Environmental Protection Agency
WHO	-	World Health Organization

CHAPTER 1

INTRODUCTION

1.1 Background of Study

It is estimated up to 2.7 million lives worldwide are potentially to be saved each year if fruits and vegetables consumption is sufficiently taken (WHO, 2013). As has been highlighted in the report of World Health Organization, most countries were encouraged to conduct health based programs to achieve the target, at least for the consumer to consume about 400 g of fruits and vegetables daily. The reasons for the recommended consumption volume basically for the prevention of several micronutrients deficiencies especially for less develop countries. For example, in United States, the *Let's Move!* Campaign (Torres, 2014) was organized by Department of Education, in conjunction with the USDA and the U. S. Department of Health and Human Services. The campaign was inspired by the fact that the obesity rate among the children in their country has increased three times, compared to over the past 30 years and their expected lifespan is now less than their parents. In 2006, New York City had launched “Move to Fruits and Vegetables” Campaign (Spear, 2007) through their Health Department and New York State Department of Agriculture and Markets. This expansion of initiative was held to increase accessibility to healthy foods amongst low income neighborhood. Similar programs had been conducted in several countries under different names, but it was held to achieve the same goal which is to increase the ease of understanding to healthy foods among low income neighborhood.

When all people in the world start focusing on staying healthy by eating fresh fruits and vegetables, some consumers may be turned down by widespread use of pesticides and high concentration of pesticides residue. Probably, some or most people had been alerted on the exposure of pesticides towards workers and farmers because they directly deal with those compounds. However, the effects of pesticides residue towards consumer were seldom discussed openly in the community discussion or social media. It became an economical issue to consumers whether to buy organic products or not. They may prefer the safer organic produce which does not use pesticides, but the cost is not affordable to lower income groups. There are few cases reported recently on the *E. coli* outbreak in California and Tennessee this year from identified its source which are from salads (Avila, 2013) and fresh cow milk (Beecher, 2013) respectively. As reported in California, from 21 patients, 2 patients were suspected with hemolytic uremic syndrome (HUS) and fatal kidney complication. Elsewhere in India for example, the Consumer Voice fought against pesticides issue at Delhi Court when they found out farmers used pesticide 750 times higher than the European standards (Arvind, 2013). In fact, some pesticides that have been used were internationally banned pesticides and commonly found in vegetables and fruits. The pesticides can cause headaches, affected fertility and even could damage kidney and liver. Besides that, scientist also found that pesticides and other poisonous chemicals that were usually used in growing fruits and vegetables can cause young children at risk of developing cancer in the future (Sanborn *et al.*, 2012).

Pesticides had increased in importance in the agricultural field since it can guarantee the growers to have a quality crop when harvesting and may extend its supermarket shelf life. Previously, organic pesticides had been popular to be used for agricultural field. However, as more pesticides being used, the need for stronger and newer synthetic pesticides keep increasing since the insects will produce new generations that are resistant to the current pesticides being used. Collaborative International Pesticides Analytical Council has categorized synthetic pesticides according to their molecular structure (CIPAC, 2009). Three groups of pesticides are

commonly used which are organophosphates, organochlorines and inorganic pesticides. Organophosphates are the most common and widely used synthetic pesticides. Compared to organochlorines and inorganic pesticides, both groups have higher toxicity and its compound break down in a shorter time period after applying it to the crops. These two pesticides; organochlorine and inorganic pesticides can cause detrimental effects on health which result in severe damage to the brain system of animals and human. Organochlorines pesticides do not easily break down like organophosphates. However, according to scientists, the application of organochlorines will cause pesticides residues to remain in the environment for a long period and can be the major cause of cancer. This group of pesticides contains toxic element such as mercury and arsenic (CIPAC, 2009). These dangerous compounds also known as persistent organic pollutants or POPs already exist for so many years in our environment. Their existence in the environment will be longer and may be incorporated into the food chain and their concentration will be magnified.

Shipman (2013) reported 48 - 81% of pesticide residues found in vegetables in the United States from the year 2000 until 2008. In Malaysia, Mazlan and Mumford (2005) had found the highest content of pesticide active ingredients in cabbage; namely indoxacarb, fipronil and chlorpyrifos combined with cypermethrin with EIQ (Environmental Impact Quotient) value of 42.97, 49.50, 52.80 and 54.50 respectively. Zawiyah *et al.* (2007) also found cypermethrin in several types of vegetables samples in the range from 0.16 to 1.48 mg/kg. Once pesticides being applied to crops, several possibilities on distribution path can be expected (Sangaralingam, 2005). Depending on their molecular structure, some of them will eventually break down after application while some of them will remain on surface of the fruits and vegetables, or remain in soil. When it rains, the rainwater will wash some of them into water supplies or groundwater which then recirculates the pesticides in the environment. Volatile pesticides that are easily evaporated will be distributed in our atmosphere and may finally return together with the raindrop. In addition, pesticides may also be distributed when contaminated fruits and vegetables were consumed. Although concrete facts and cases on pesticides

poisoning cannot be confirmed, the possible danger and effect just cannot be ignored. The acute effect will be through accumulation in our body system and can lead to death. In this study, the used of pesticide focused on eight of organophosphorus pesticides because most of them were listed as common pesticide that can be found in vegetables.

Chen *et al.* (2013) suggested ozone treatment for handling hydrophobic pesticides. This method can be considered as innovative technique but the cost to build the instrument will be of economic issues. Other researcher also had tried proposing hydrostatic pressure treatment as one of the alternatives (Iizuka *et al.*, 2013 (Part 1); Iizuka *et al.*, 2013 (Part 2)). But again, the same factor which is cost effective needs to be considered. Zohair had developed analysis of different washing solution to enhance the washing off effect of several pesticides (Zohair, 2001). The solutions involved varied from acidic, neutral and alkaline solution as well as tap water. He found that acidic washing solution is better for organochlorine pesticides removal as compared to tap water. Overall, it can be concluded that most study of pesticide removal need addition of detergent instead of using only tap water (Liang *et al.*, 2012; Fernandez *et al.*, 2013).

Pesticides were commonly found in vegetables and fruits since its application were important as protection from insects. Pesticides removal technique were proposed in several studies as mentioned earlier which has their various advantages and disadvantages depending on the consumer's condition. Apart from pesticide, heavy metal was also one of the major problems in food business. This problem had been studied by Azelee *et al.* (2013) who proposed heavy metals removal like arsenic (As), lead (Pb), cadmium (Cd), nickel (Ni) and mercury (Hg) from *Perna viridis* using chelation technique. The removal performance obtained were 54.34% (As), 96.79% (Hg), 82.89% (Hg), 75.02% (Cd) and 61.83% (Ni). Mussels are used as biological indicator (Bedford *et al.*, 1966) because of the capabilities as filter feeder. Mussels were also found to have various organic compounds (Somchit *et al.*, 2009) and pesticides (Sivalingam, 1982). Since heavy metals were successfully removed using the chelation

technique, the removal potential study of pesticides using the same technique maybe viable.

Pesticides contamination had been seen in various food products as mention earlier especially in vegetables. Besides pesticides residue, heavy metal also had reported as one of the contaminant in vegetables (Garcia *et al.*, 1974; Fang *et al.*, 2014; Kang *et al.*, 2008 and Kanakaraju *et al.*, 2007). Unknowingly, all these studies showed that our food products especially vegetables contain various contaminants which lead to consumer's health problems. Therefore, in this study, a consumer product in market which is "Fruit and Vegie Wash" from Australia was tested on its effectiveness in removing pesticides and heavy metals by using method which had been proposed by Wang *et al.* (2013) and Azelee *et al.* (2013).

1.2 Problem of Statement

Nowadays, consumers are moving towards adopting a healthy lifestyle including making sure all the food that they eat is not contaminated. Washing using tap water is the most commonly used method in our community. However, as reported by Geetanjali *et al.* (2009), washing by using various type of solution had gave difference results in term of dissipation percentage of pesticides level. Although the types of pesticides removed were same which are hexachlorobenzene (HCB), lindane, p,p-dichlorodiphenyltrichloroethane (p,p-DDT), dimethoate, profenofos and pirimiphos-methyl, but this study had shown that the percentage of pesticides dissipation can be increased by adding acetic acid or sodium chloride into tap water. Table 1.1 below shows the percentage of pesticides residue dissipation review of different washing solutions towards tomatoes sample.

Table 1.1: Review on percentage of residue dissipation (Geetanjali *et al.*, 2009).

Processing	Commodity	Pesticide	Initial residues (ppm)	% Residue dissipation
Washing Acetic acid solution	Tomatoes	HCB	1	51.3
		Lindane	1	47.0
		P,p-DDT	1	33.7
		Dimethoate	1	91.5
		Profenofos	1	86.0
		Pirimiphos-methyl	1	93.7
		HCB	1	42.9
		Lindane	1	46.1
		P,p-DDT	1	27.2
		Dimethoate	1	90.8
Sodium chloride (at 10% NaCl)		Profenofos	1	82.4
		Pirimiphos-methyl	1	91.4
		HCB	1	9.62
		Lindane	1	15.3
		P,p-DDT	1	9.17
		Dimethoate	1	18.8
		Profenofos	1	22.7
		Pirimiphos-methyl	1	16.2
Tap water		HCB	1	9.62
		Lindane	1	15.3
		P,p-DDT	1	9.17
		Dimethoate	1	18.8
		Profenofos	1	22.7
		Pirimiphos-methyl	1	16.2

In Malaysia, a commercial consumer product from Australia claimed to be effective in removing pesticides. This product was available in several hypermarkets in Malaysia such as Isetan and Jaya Jusco Supermarket. However, the response from consumers was not encouraging. This situation was probably caused by the low level of awareness from Malaysian citizens on the culture of healthy lifestyle, expensive price and insufficient promotion from the manufacturer.

Other than current commercial products in the markets, there are many tips or homemade recipes which have been shared through social medium. They used blogs, website and YouTube as medium to share their recipes on preparation of washing solution to remove any harmful bacteria and also pesticides residue. Usually, the

ingredients used are common kitchen goods such as lemon juice, baking soda, vinegar and salt. These practices have gain lots of positive feedback from consumer since the cost is cheaper and the formulation is easy to be prepared. However, the effectiveness of this technique does not proven scientifically since no study had been carried out based on these homemade recipes. These practices can be nurtured among Malaysian citizens since it may give benefits towards a healthier life. Unknowingly, most population may be exposed to various pesticides through consumption of fruits and vegetables, the pesticide residues tend to accumulate in the body. Concerns have been raised about the possible role of continuous low-dosage exposure in causing certain cancers.

The study on removal of heavy metals from mussel samples have been successfully carried out in 2013 (Azelee *et al.*, 2013). The successful chelation technique was tested to study the possibility of removing pesticides from food products such as vegetables and mussels. Finally, this study was further continued to develop combine methods to analyse the performance of pesticide and heavy metals removal either using both methods simultaneously or through continuous washing process and chelation technique.

1.3 Objectives of Study

The objectives of the study are as follows:

- i. To determine percentage reduction of pesticides in vegetable and mussel samples using commercial washing solution through washing treatment.
- ii. To optimize the washing and chelating treatment conditions for optimum removal of pesticides and heavy metals.
- iii. To evaluate levels of pesticide and heavy metal residues via the simultaneous and continuous of commercial washing solution and sodium acetate chelating solution.

1.4 Scope of Study

Washing technique was applied to long beans, broccoli, lettuce, cucumber, tomatoes and mussel samples. The selection of various vegetables is based on different types of skin surface and among the vegetables that are often eaten as raw or made into salad. Mussels represent food sample from marine group. Azinphos-methyl, chlorpyrifos, dichlorvos, disulfoton, ethoprophos, fenchlorphos, parathion-methyl, prothiofos and malathion are chosen for this study because they were listed as the frequent pesticides used by the farmers in Malaysia (DOA, 2014). Removal of metals include arsenic (As), lead (Pb), cadmium (Cd) and nickel (Ni) which are the common toxic heavy metals by using ammonium acetate as chelating solution. Several normal home practices were optimized during washing technique with focus on washing solution, washing time, temperature and swirl effect to achieve maximum removal rate of pesticides. The extract of sample for pesticides was analysed using gas chromatography that equipped with electron capture detector (GC-ECD) whereas analysis of metal using inductively coupled plasma optical emission spectrometry (ICP-OES). Simultaneous removal of pesticides and heavy metal was studied using a proposed design with the expected percentage of removal performances.

In previous studies, many techniques have been proposed as pesticide and heavy metals residues removal. Some of the studied parameters in previous studies of washing were evaluated in this study including time, volume, temperature and swirl effect. However, ultrasonic effect was not included in this study because ultrasonic equipment is rare to be found in normal household appliances. This study was developed according to normal and common practices of consumer during the preparation of food.

1.5 Significant of Study

Compared to hydrostatic treatment, ozone treatment or washing with tap water, washing with commercial solution provides several benefits. This technique is simple with better performance in removing pesticides. It is practical to be done by users without specific skills or equipment. Washing technique maintains the freshness of food products without any adverse effect to human health. Besides that, this study provided a scientific explanation on the suitable conditions for washing method to achieve maximum rate of pesticides and heavy metals removal. This study is hoped to provide assurance for consumer rather than became just as traditional remedies only. In fact, this study can help in exporting market of mussels and other food product, since the level of pesticides and heavy metals can conform to the standard for human food consumption.

From the economic view, the commercial product used in this study provides several advantages to consumers. The market price for this product in Malaysia is about RM 40.00 for a bottle of 750 mL of concentrated washing solution. Normal volume per usage is about 5 mL in 1000 mL of tap water. It can be estimated that one bottle of this commercial product can maximally use for about 150 times which equal to about 5 months. Consumer only spending their money about RM 0.27 for every daily use. In comparison, consumer will face various prices if they prefer in using homemade remedies such as vinegar (RM 1.99/330 mL), salt (RM 1.90/800gm) and lemon (RM 1.50/pcs). This study purposely developed to satisfy and meet the needs of busy consumers, particularly for users with careers because this product is easily to be used by only mixing with tap water. This comparison of economic price is for usage with soaking style. If the consumer prefer in washing with spraying style, a bottle of this commercial product will be last for more than 5 months. Hence, more benefits are gain by the consumer.

REFERENCES

- Abou-Arab, A.A.K. (1999). Behavior of Pesticides in Tomatoes during Commercial and Home Preparation. *Food Chemistry*. **65**, 509-514.
- Adachi, A. and Okano, T. (2006). Pesticide Residue Reduction in Selected Vegetables Using Rice-Bran. *Journal Health Science*, **42** (3):320-323.
- Alam, M. G. M. and Snow, E.T. and Tanaka, A. (2003). Arsenic and Heavy Metal Contamination of Vegetables Grown in Samta Village, Bangladesh. *Science of the Total Environment*. **308** (1–3), 83–96.
- Aros, N. M. T. (1985). Pesticide Poisoning Studies and Data Collection in Malaysia. Toxicologist. Pesticides Section, Department of Agriculture Malaysia.
- Arvind, A. (2013). Article of High Court Rejects Delhi Government's Pesticides Report as Ngo Accuses State of 'Whitewashing' The Threat to Capital's Fruit and Vegetables. Mail Online India.
- Avila, W. (2013). Article of California Company Recalls Salads, Wraps Linked to E. Coli Outbreak. NBC Southern California.
- Aweng, E. R., Karimah, M. and Suhaimi, O. (2011). Heavy Metals Concentration of Irrigation Water, Soils and Fruit Vegetables in Kota Bharu Area, Kelantan, Malaysia. *Journal Applied Sciences in Environment Sanitation*, **6**(4), 463-470.
- Azelee, I. W. Catalytic Chelation Technique for the Removal of Toxic and Heavy Metals from Green Mussel, *Perna Viridis*. Degree of Master of Science (Chemistry) Thesis. Universiti Teknologi Malaysia; 2013.
- Babji, A. S., Embong, M. S. and Woon, W. W. (1979). Heavy Metal Contents in Coastal Water Fishes of West Malaysia. Bull. *Journal Environment Contamination Toxicology* **23**, 830-836.

- Bedford, J. M., Roelof, E. W. and Zabik, M. J. (1966). The Freshwater Mussel as a Biological Monitor of Pesticide Concentrations in a Lotic Environment. M. S. Degree. Faculty of Michigan State University, East Lansing 48823.
- Beecher, C. (2013). Article of Tennessee *E. Coli* Outbreak Linked to Raw Milk Dairy. Food Safety News. United States.
- Biopharmaceutics. (2003). *Guidance for Industry; Bioanalytical Method Validation*. Revision 1. U.S. Department of Health and Human Services.
- Carson, R. (2015, August 13). The Story of Silent Spring, Natural Resources Defense Council. Retrieved January 9, 2016, from <http://www.nrde.org/stories/story-silent-spring>.
- Chai, L. K. and Elie, F. (2009). A Rapid Multi-residue Method for Pesticide Residues Determination in White and Black Pepper (*Piper Pigrum L.*). *Journal Food Control*. **32**, 322-326.
- Chen, J. Y., Lin, Y. J. and Kuo, W. C. (2013). Pesticide Residue Removal from Vegetables by Ozonation. *Journal Food Engineering*. **114**, 404–411.
- Cieslinski, G., Van Rees, K. C. J., Szmigielska, A. M., Krishnamurti, G. S. R. and Huang, P. M. (1998). Low-Molecular-Weight Organic Acids in Rhizosphere Soils of Durum Wheat and their Effect on Cadmium Bioaccumulation. *Journal Plant and Soil*. **203** (1), 109-117.
- Clark, J. (2004, May 15). Introducing Esters. Chemical Guide U. K. Retrieved January 18, 2016, from www.chemguide.co.uk/organicprops/ester/background.html.
- Colin Baird. (2000). Environmental Chemistry, W. H. Freeman and Company, New York.
- Collaborative International Pesticides Analytical Council (CIPAC, 2009). MT Methods. www.cipac.org.
- Commission of the European Communities (EU). (2004). Monitoring of Pesticide Residues in Products of Plant Origin in the European Union, Norway, Iceland and Liechtenstein (Part 1). SEC1416. Retrieved January 18, 2016 from http://ec.europa.eu/food/fvo/specialreports/pesticide_residues/report_2004_en.pdf.
- Department of Agriculture Malaysia (DOA). (2014, February 1). Perangkaan Tanaman. Retrieved December 6, 2015 from <http://www.doa.gov.my/web/guest/data-perangkaan-tanaman>.

- Di Maio, V. J. M. (2001). *Forensic Pathology*, 2nd ed., CRC Press, Boca Raton, FL.
- Drum, C. (1980). *Soil Chemistry of Pesticides*, PPG Industries, Inc. USA.
- Engineering Services Division, Ministry of Health Malaysia. (2000). National Standard for Drinking Water Quality. Kementerian Kesihatan Malaysia, D1.
- Fang, B. and Zhu, X. (2014). High Content of Five Heavy Metals in Four Fruits: Evidence from a Case Study of Pujiang County, Zhejiang Province, China. *Journal Food Control*. **39**, 62-67.
- Fernandez, O. L., Otero, R. R. and Gándara, J. S. (2013). Factors Governing the Removal of Mancozeb Residues from Lettuces with Washing Solutions. *Journal Food Control*. **34**, 530-538.
- Food Safety and Quality Division. (2000). *Food Safety Standards in Major Export Markets: A Readymade Guide for Agro Exporters*. [Brochure]. Ministry of Health: Malaysia.
- Garcia, W. J., Blessin, C. W. Inglett, G. E. and Carlson, R. O. (1974). Physical-Chemical Characteristics and Heavy Metal Content of Corn Grown on Sludge-Treated Strip-Mine Soil. *Journal Agriculture Food Chemistry*. **22**(5), 810-815.
- Geetanjali, K., Santosh, S. and Naik, S. N. (2009). Food Processing a Tool to Pesticide Residue Dissipation – A review. *Journal Food Research International*. **42**, 26-40.
- Graziela, C. R. M., Andrade, Monteiro, S. H., Francisco, J. G., Figueiredo, L. A., Rocha, A. A. and Tornisielo, V. L. (2015). Effects of Types of Washing and Peeling in Relation to Pesticides Residues in Tomatoes. *Journal Brazilian Chemical Society*. 26(10), 1678-4790.
- Greene, B. (1984). Removal of Heavy Metal Ions from Contaminated Water by *Chlorella Vulgaris*. Doctoral Degree of Chemistry Thesis. New Mexico State University.
- Heber, D. Vegetables, Fruits and Phytoestrogens in the Prevention of Diseases. (2004). *Journal Postgraduate Medicine*. **50**, 145–149.
- Herdman, R.C., Pahrham, W.E. and Shen, S. Pesticide Residues in Food: Technologies for Detection. Government Printing Office, Washington (1988).
- Hizal, J. Modeling of Heavy Metal Adsorption onto Clay Minerals. Ph.D. Thesis, Istanbul University, Turkey; 2001.

- Hizal, D. B. Adsorption of Olive Leaf Antioxidants on Silk Fibroin. Degree of Master of Science (Chemical Engineering) Thesis. Izmir Institute of Technology; 2006.
- Iizuka, T., Maeda, S. and Shimizu, A. (2013). Removal of Pesticide Residue in Cherry Tomato by Hydrostatic Pressure (Part 1). *Journal Food Engineering*. **116**, 796–800.
- Iizuka, T. and Shimizu, A. (2013). Removal of Pesticide Residue in Cherry Tomato by Hydrostatic Pressure (Part 2). *Journal Innovative Food Science and Emerging Technologies* INNFOO-01005, 6-11.
- International Conference on Harmonisation (ICH). (1996). *Validation of Analytical Procedures: Text and Methodology*. International Conference on Harmonisation of Technical Requirements for Registration of Pharmaceuticals for Human Use.
- Isa, I. M., Wardani, N. I., Hashim, N. and Ghani, S. A. (2015). Determination of Trace Level Copper (II) in Malaysian Vegetables by Cyclic Voltammetry. *International Journal Electrochemical Science*. **10**, 498-503.
- Islam, M. R., Salminen, R. and Lahermo, P. W. (2000). Arsenic and Other Toxic Elemental Contamination of Groundwater, Surface Water and Soil in Bangladesh and its Possible Effects on Human Health. *Journal Environmental Geochemistry Health*. **22**, 33–53.
- Islam, M. S. and Hoque, M. F. (2014). Concentrations of Heavy Metals in Vegetables around the Industrial Area of Dhaka city, Bangladesh and Health Risk Assessment. *International Food Research Journal*. **21**(6), 2121-2126.
- Ismail, I. and Saleh, I. M. (2012). Analysis of Heavy Metals in Water and Fish (Tilapia Sp.) Samples from Tasik Mutiara, Puchong. *Malaysian Journal of Analytical Sciences*. **16**(3), 346-352.
- Jaafar, M. H. (2010). Determination of Heavy Metal Accumulation in Ipomoea Aquatic using Wet Destruction Method. Degree of Bachelor of Chemical Engineering.
- Jalal, K. C. A., John, A. Habab, M., Mohd, A. Y. and Kamaruzzaman, B. Y. Bioaccumulation of Selected Metals in Fresh Water Haruan Fish (*Channa Striatus*) Collected from Pahang River Basin, Malaysia. *Oriental Journal Chemistry*. **29**(4), 1553-1558.
- Jarup, L. (2003). Hazards of Heavy Metal Contamination. *Oxford Journals*. **68** (1), 167-182.
- Jlbris. (2014, February 23). Aloe vera. School of Chemistry, University of Bristol. Retrieved December 20, 2015 from <http://chempics.wordpress.com/category/aloe-vera>.

- Kamaruzzaman, B. Y., Ong, M. C., Zaleha, K., Shahbudin, S. (2008). Levels of Heavy Metals in Green-Lipped Mussel *Perna veridis* (Linnaeus) from Muar Estuary, Johore, Malaysia. *Pakistan Journal Biological Sciences*. **11**, 2249-2253.
- Kanakaraju, D., Mazura, N. A. and Khairulanwar, A. (2007). Relationship between Metals in Vegetables with Soils in Farmlands of Kuching, Sarawak. *Malaysian Journal of Soil Science*. **11**, 57-69.
- Kang, J. H., Park, H., Chang, Y. S. and Choi, J. W. (2008). Distribution of Organochlorine Pesticides (OCPs) and Polychlorinated Biphenyls (PCBs) in Human Serum from Urban Areas in Korea. **73** (10), 1625–1631.
- Kasiotis, K. M., Emmanouil, C., Anastasiadou, P., Papadi-Psyllou, A., Papadopoulos, A. Okay, O. Machera, K. (2015). Organic Pollution and Its Effects in the Marine Mussel *Mytilus galloprovincialis* in Eastern Mediterranean Coasts. *Chemosphere*. **119**, 145–152.
- Khairiah J., Zalifah, M. K., Yin, Y. H. and Aminah, A. (2004). The Uptake of Heavy Metals by Fruit Type Vegetables Grown in Selected Agricultural Areas. *Pakistan Journal of Biological Sciences*. **7** (8), 1438-1442.
- Kin, C. M. and Huat, T. G. (2010). Headspace Solid-phase Microextraction for the Evaluation of Pesticide Residue Contents in Cucumber and Strawberry after Washing Treatment. *Food Chemistry*. **123**, 760–764.
- Koduru, J. R., Chang, Y. Y. and Kim, I. S. (2014). Low-Cost Schizandra Chinesis Fruit Peel for Co (II) Removal from Aqueous Environment: Adsorption Properties and Mechanism. *Asian J. of Chem*. 26(1), 289-297.
- Krishnamurthy, S. (1992). Biomethylation and Environmental Transport of Metals. *Journal Chemical Education*. **69**(5), 347.
- Lembaga Racun Makhluks Perosak (2009, May 1). Senarai Racun Makhluks Perosak Berdaftar. Portal Rasmi Jabatan Pertanian. Retrieved July 1, 2014, from www.doa.gov.my.
- Liang, Y., Wang, W., Shen, Y., Liu, Y. and Liu, X. J. (2012). Effects of Home Preparation on Organophosphorus Pesticide Residues in Raw Cucumber. *Food Chemistry*. **133**, 636–640.

- Lindaverrill, Lando, A. M. and O' Connell, K. (2012). Consumer Vegetable and Fruit Washing Practices in the United States 2006 and 2010. *Journal Food Protection Trends*. **32**(4), 164-172.
- Li, X. and Thornton, I. (2001). Chemical Partitioning of Trace and Major Elements in Soils Contaminated by Mining and Smelting Activities. *Journal Applied Geochemistry*. **16**, 1693–1706.
- Lukasik, J., Bradley, M. L., Scott, T. M., De, M., Koo, A., Hsu, W. Y., Bartz, J. A. and Farrah, S. R. (2003). Reduction of Poliovirus, Bacteriophages, *Salmonella Montevideo*, and *Escherichia Coli* O157:H7 on Strawberries by Physical and Disinfectant Washes. *Journal Food Protection*. **66**(2), 188-193.
- Madivoli, E. S., Gitu, L. and Gumba, E. (2012). Isolation and Identification of Essential Oils from *Cymbopogon Citratus* (STAPF) Dc Using GC-MS and FT-IR. *Journal Chemical and Mathematics Research*, **2**(4), 13-22.
- Majewski, M. S., Zamora, C., Foreman, W. T. and Kratzer, C. R. (2006). Contribution of Atmospheric Deposition of Pesticide Loads in Surface Water Runoff. U. S. Geological Survey Report. 2005-1307.
- Mazlan, N. and Mumford, J. (2005). Insecticide Use in Cabbage Pest Management in the Cameron Highlands, Malaysia. *Journal Crop Protection*. **24**, 31–39.
- National Academy of Sciences. (1968). Effects of Pesticides on Fruit and Vegetable Physiology. eBook of Principles of Plant and Animal Pest Control. Library of Congress Catalog Card Number 68-60085. **6**, 51-55.
- Nownes, A.J. (1991). Interest Groups and the Regulation of Pesticides: Congress, Coalitions and Closure. *Journal Policy Sciences*. **24**, 1-18.
- Peijnenburg, W. J. G. M. and Jager, T. (2003). Monitoring Approaches to Assess Bioaccessibility and Bioavailability of Metals: Matrix Issues. *Journal Ecotoxicology Environmental Safety*. **56** (1), 63–77.
- Pesticides Act 1974. (2006). The Commissioner of Law Revision, Malaysia Under the Authority of the Revision of Laws Act 1968.
- Pena, A. G., Franeschi, F. A., Estrada, M. C., Ramos, V. M., Zarracino, R. G., Zavala Loria, J. C. and Cordova Quiroz, A. V. (2014). Fourier Transform Infrared-Attenuated Total

- Reflectance (FTIR-ATR) Spectroscopy and Chemometric Techniques for the Determination of Adulteration in Petrodiesel/Biodiesel Blends. *Quimica Nova*, **37**(3), 392-397.
- Ping, Z., Huiling, Z. and Wensheng, S. (2009). Biotransfer of Heavy Metals along a Soil-Plant-Insect-Chicken Food Chain: Field study. *Journal Environment Sciences*. **21**, 849–853.
- Racke, K. D., Steele, K. P., Yoder, R. N., Dick, W. A. and Avidov, E. (1996). Factors Affecting the Hydrolytic Degradation of Chlorpyrifos in Soil. *Journal Agricultural Food Chemistry*. **44** (6), 1582–1592.
- Ray, A., Gupta, S. D. and Ghosh, S. (2013). Evaluation of Anti-oxidative Activity and UV Absorption Potential of the Extracts of Aloe Vera L. Gel from Different Growth Periods of Plants. *Industrial Crops and Products*. **49**, 712- 719.
- Report of the Workshop on Globally Harmonised System for Malaysia Government Agencies. (2006). Department of Occupational Safety and Health Ministry of Human Resources Malaysia.
- Rintos. (2014, March 16). Pesticides-a double-edged sword. Borneo Post Online, Retrieved November 4, 2014, from www.theborneopost.com.
- Sanborn, M., Bassil, K., Vakil, C., Kerr, K. and Ragan, K. (2012). Systematic Review of Pesticide Health Effects. Ontario College of Family Physicians (OCFP).
- Sangaralingam, M. (2005). International POPs Elimination Project (IPEP). Malaysia Country Situation Report.
- Sarman, S., Hashim, N. and Kantasamy, N. (2006). A Comparative Market Basket Study of Certain Heavy Metals in Selected Vegetables. University Teknologi MARA.
- Serrano, M., Lin, A. W., McCurrach, M. E., Beach, D. and Lowe, S. W. (1997). Oncogenic Ras Provokes Premature Cell Senescence Associated with Accumulation of p53 and p16INK4a. *National Center for Biotechnology Information, U.S. National Library of Medicine*. **88** (5), 593-602.
- Shah, J., Jan, M. R., Shah, Z. and Ishrat, S. (2001). Effect of Pesticides on Soil Microorganisms, American Laboratory. **33**(23), 24.

- Shipman, D. R. (2013). Pesticide Data Program. Annual Summary, Calendar Year 2011. Department of Agriculture, United States.
- Shokrzadeh, M. and Saravi, S. S. S. (2000). Pesticides in Agricultural Products: Analysis, Reduction, Prevention. University of Medical Sciences, Sari Iran.
www.intechopen.com.
- Sivalingam, P. M. (1982). Mussel Watch of PCBs and Persistent Pesticide Residues in *Perna viridis Linnaeus* from Malaysian and Singapore Waters. *Journal Applied Geophysics*. **2**, 231-237.
- Somchit, N., Somchit, M. N., Hadi, S. A. and Zakaria, M. P. (2009). Persistent Organic Chemicals in Malaysian Waters: A Review. *Research Journal of Environmental Toxicology*. **3** (2):101-112.
- Soylak, M., Tuzen, M., Narin, I. and Sari, H. (2004). Comparison of Microwave, Dry and Wet Digestion Procedures for the Determination of Trace Metal Contents in Spice Samples Produced in Turkey. *Journal Food and Drug Analysis*. **12** (3), 254-258.
- Spear B.A. (2007). NCR for Health and Safety in Child Care and Early Education.
- Sze, P. W. C. and Lee, S. Y. (1995). The Potential Role of mucus in the depuration of copper from the mussels *Perna viridis* (L.) and *Septifer virgatus* (Wiegmann). *Mar. Pollut. Bull.* **31**, 390-393.
- Torres, A. (2014, February 3). Childhood Obesity has Declined, but that's no Thanks to the First Lady's Initiative. National Review. Retrieved September 11, 2014, from <http://www.nationalreview.com>.
- Wang, Z., Huang, J., Chen, J. and Li, F. (2013). Effectiveness of Dishwashing Liquids in Removing Chlorothalonil and Chlorpyrifos Residues from Cherry Tomatoes. *Chemosphere*. **92**, 1022-1028.
- World Health Organization (WHO). (1989). Guidelines for Predicting Dietary Intake of Pesticide Residues, World Health Organization, Geneva. 24.
- WHO Technical Report Series. (2013). Diet, Nutrition and the Prevention of Chronic Diseases. Report of a Joint FAO/WHO Expert Consultation Geneva, World Health Organization, No. 916.

- Xu, X., Cao, X. and Zhao, L. (2013). Comparison of Rice Husk- and Dairy Manure-Derived Biochars for Simultaneously Removing Heavy Metals from Aqueous Solutions: Role of Mineral Components in Biochars. *Chemosphere*. **92** (8), 955–961.
- Yuan, G. F., Sun, B., Yuan, J. and Wang, Q. M. (2009). Effects of Different Cooking Methods on Health-Promoting Compounds of Broccoli. *Journal Zhejiang University Science*. **10** (8), 580–588.
- Zainudin, B. H., Salleh, S. Mohamed, R., Yap, K. C. and Muhamad, H. (2015). Development, Validation and Determination of Multiclass Pesticide Residues in Cocoa Beans using Gas Chromatography and Liquid Chromatography Tandem Mass Spectrometry. *Journal Food Chemistry*. **172**, 585–595.
- Zacharia and Tano, J. (2011). Identity, Physical and Chemical Properties of Pesticides. University of Dar es Salaam, Tanzania. www.intechopen.com.
- Zawiyah, S., Che Man, Y. B., Nazimah, S. A. H., Chin, C. K., Tsukamoto, I., Hamanyza, A. H. and Norhaizan, I. (2007). Determination of Organochlorine and Pyrethroid Pesticides in Fruit and Vegetables using SAX/PSA Clean-up Column. *Food Chemistry*. **102**, 98–103.
- Zhang, Z. Y., Liu, X. J. and Hong, X. Y. (2007). Effects of Home Preparation on Pesticide Residues in Cabbage. *Journal Food Control*. **18**, 484–487.
- Zhuang, P., McBride, M. B., Xia, H., Lia, H. and Lia, Z. (2009). Health Risk from Heavy Metals via Consumption of Food Crops in the Vicinity of Dabaoshan mine, South China. *Science of the Total Environment*. **407**, 1551–1561.
- Zohair, A. (2001). Behaviour of Some Organophosphorus and Organochlorine Pesticides in Potatoes during Soaking in Different Solutions. *Journal Food and Chemical Toxicology*. **39**, 751–755.